

The Main Characteristics of the Wills-Harrison Effective Pair Potential in Liquid Co

Natalia D. Vatolina

Ural Federal University, Mira st. 19, 620002 Ekaterinburg, Russia
ndvatolina@gmail.com

Copyright © 2013 Natalia D. Vatolina. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

It is found that the position of the first minimum of the Wills-Harrison effective pair potential in liquid Co begins to shift significantly to the right-hand side when the d - d -non-diagonal coupling begins to predominate in the metal under consideration.

Keywords: Transition metal, Wills-Harrison pair potential, d -state coupling

In [1] the Wills-Harrison (WH) model [2] was corrected by means the introduction the probability p that all 25 d - d couplings between two different atoms are equiprobable and probability $(1 - p)$ that only 5 equiprobable diagonal couplings are possible.

Here, we consider how the magnitude p influences the position, r_{\min} , and the magnitude of the first minimum of the WH effective pair potential, $\varphi_{\text{WH}}(r)$, in liquid Co. Earlier, the analogous study was fulfilled for liquid Fe [3].

We use the local Bretonnet-Silbert (BS) model pseudopotential [4] for description the s -electron contribution to $\varphi_{\text{WH}}(r)$. Input data (WH and BS parameters and the experimental mean atomic volume, Ω) are listed in Table 1.

It is found that behaviors of both curves (Figs. 1- 2) are similar that in liquid Fe [3]: the magnitude of the first minimum quite monotonously increases with increasing p , but the first-minimum position begins to increase harshly from $p = 0.6$ approximately. It denotes that significant shift of the $\varphi_{\text{WH}}(r)$ first minimum to the right-hand side is occurred in liquid Co when the non-diagonal coupling begins to predominate. The same tendency was found in liquid Fe [3].

Table 1. Input data for calculation

r_d (a.u.) [2]	z_s [5]	z_d [5]	R_C (a.u.) [5]	a (a.u.) [5]	Ω (a.u.) [6]
1.437	1.4	7.6	1.64	0.393	85.85




Figure 1. r_{\min} of $\varphi_{\text{WH}}(r)$ in liquid Co at different p ($T=1863\text{K}$).




Figure 2. $\varphi_{\text{WH}}(r_{\min})$ in liquid Co at different p ($T=1863\text{K}$).

References

- [1] N.E. Dubinin, Account of non-diagonal coupling between d electrons at describing the transition-metal pair potentials, *J. Phys.: Conf. Series*, 338 (2012), 012004.
- [2] J.M. Wills, W.A. Harrison, Interionic interactions in transition metals, *Phys. Rev. B*, 28 (1983), 4363-4373.
- [3] N.D. Vatolina, The main characteristics of the Wills-Harrison effective pair potential in liquid Fe, *Adv. Studies Theor. Phys.*, 7 (2013), in print.
- [4] J.L. Bretonnet, M. Silbert, Interionic interactions in transition metals. Application to vanadium, *Phys. Chem. Liq.*, 24 (1992), 169-176.
- [5] N. Jakse, J.L. Bretonnet, Structure and thermodynamics of liquid transition metals: integral-equation study of Fe, Co and Ni, *J. Phys.: Condens. Matter*, 7 (1995), 3803-3815.
- [6] Y. Waseda, *The Structure of Non-crystalline Materials – Liquids and Amorphous Solids*, McGraw-Hill, New York, 1981.

Received: June 28, 2013